



**UNIVERSITI PUTRA MALAYSIA**

**MICROPROPAGATION OF MANGOSTEEN (*GARCINIA  
MANGOSTANA* L.) BY DIRECT AND INDIRECT ORGANOGENESIS**

**SUCI RAHAYU**

**FP 2007 3**

**MICROPROPAGATION OF MANGOSTEEN  
(*GARCINIA MANGOSTANA* L.) BY DIRECT AND  
INDIRECT ORGANOGENESIS**

**SUCI RAHAYU**

**MASTER OF SCIENCE  
UNIVERSITI PUTRA MALAYSIA**

**2007**



**MICROPROPAGATION OF MANGOSTEEN (*GARCINIA MANGOSTANA*  
L.) BY DIRECT AND INDIRECT ORGANOGENESIS**

**By**

**SUCI RAHAYU**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Master of Science**

**April 2007**



## **DEDICATION**

I dedicate this thesis to my father, Gito Soewarno and my mother, Subariah.

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirement for the degree of Master of Science

**MICROPOPAGATION OF MANGOSTEEN (*Garcinia mangostana* L.) BY  
DIRECT AND INDIRECT ORGANOGENESIS**

By

**SUCI RAHAYU**

**April 2007**

**Chairman : Associate Professor Mhdzar Abdul Kadir, PhD**

**Faculty : Agriculture**

The mangosteen (*Garcinia mangostana* L.) still remains an under exploited tropical fruit species that has potential to be developed for local consumption and export commodity. The problem for large scale planting of this plant is not having enough planting materials because the conventional vegetative propagation method has low percentage of success. One of the alternative approaches for mass propagation of mangosteen is through tissue culture. The advantages of propagation through tissue culture are mass propagation can be achieved in a short of time period, plantlet can be produced all year round, and require less space of the plant in the laboratory compare with to the field. The result of the study on *in vitro* propagation of mangosteen in terms of production of adventitious shoots through direct organogenesis and indirect organogenesis in term of callus induction was conducted at the laboratory of Faculty of

Agriculture, Universiti Putra Malaysia. It was observed that among 24 initiation media tested, 16 mg/l BAP gave the highest number of shoots (7.00 shoot/explant), followed by 8 mg/l BAP (6.3 shoots/explant) after eighth weeks of culture. In general, the supplement of AdS in media did not significantly influence the number of shoots produced.

There was negative correlation between the height of shoot and the number of shoot produced. The highest number of shoots was recorded on media with 1 mg/l and 2 mg/l BAP, while the shortest shoot was on media of 16 mg/l and 8 mg/l BAP. Contrary to the height of shoot, there was positive correlation between the weight of shoots and the number of shoots. Treatment with 16 mg/l and 8 mg/l BAP gave the highest weight of shoots produced per seed explant.

During subculture, treatment with BAP 2 mg/l gave the highest number of shoots at every occasion of subculture (39.56 shoots per explant), followed by 1 mg/l BAP (27.71 shoots per explant). Height of shoots on all media of subculture did not significantly differ. Combination of 1 mg/l BAP and 2 mg/l Kinetin was the most suitable media for elongation of the shoot (1.06 cm).

Among the rooting media,  $\frac{1}{2}$  MS + 15 mg/l NAA + 0.05% Charcoal + 10 g/l sucrose was most suitable for the emergence of roots with 3.4 roots per shoot). While  $\frac{1}{2}$  MS + 1 mg/l IAA + 0.05% Charcoal + 10 mg/l sucrose seemed to be most suitable media for lengthening the roots. The average length of root on this

media was 6.85 cm per shoot. Plantlets from rooting experiments acclimatized on the media of soil + sand + peat (2 : 1: 1) have grown well. 100% of them could survive under misting chamber condition for eight weeks.

In study on organogenesis through callus induction, all media with the exception of MS, BAP alone and 2,4-D alone could produce calli.

It is expected that this findings can contribute to the knowledge of science, particularly on *in vitro* propagation of mangosteen, and can be used as a reference for other students and researchers as well as agricultural industries in developing mass propagation of mangosteen.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PEMBIAKAN MIKRO MANGGIS (*Garcinia mangostana* L.) MELALUI  
ORGANOGENESIS SECARA LANGSUNG DAN TAK LANGSUNG**

Oleh

**SUCI RAHAYU**

**April 2007**

**Pengerusi : Profesor Madya Mihdzar Abdul Kadir, PhD**

**Fakulti : Pertanian**

Manggis (*Garcinia mangostana*) merupakan antara spesies buah tropika yang belum diterokai, meskipun ianya mempunyai potensi untuk dikembangkan sama ada untuk pemakanan tempatan mahupun untuk komoditi eksport. Permasalahan yang berlaku adalah tidak cukupnya anak pokok untuk penanaman skala besar. Pembiakan vegetatif secara biasa mempunyai peratus kejayaan yang rendah. Salah satu pilihan untuk pembiakan secara besar-besaran ialah dengan kultur tisu. Keuntungan pengembang biakan dengan kultur tisu adalah pembiakan secara besar-besaran dapat diperolehi dalam waktu yang lebih pendek, produksi anak pokok dapat diperolehi sepanjang tahun dan jimat ruang penyimpanan dalam makmal berbanding penyimpanan anak pokok di lapang. Hasil kajian dari penyelidikan kultur tisu dengan tajuk Pembiakan Tampang Manggis Melalui Organogenesis Secara Langsung dan Tidak langsung (Induksi Kalus) adalah



sebagai berikut: Diantara 24 media inisiasi yang dikaji, 16 mg/l BAP memberikan jumlah tunas yang paling banyak, iaitu 7.00 tunas tiap eksplan, diikuti oleh 8 mg/l BAP yang menghasilkan 6.3 tunas selepas lapan minggu penanaman di media. Secara amnya, pencampuran AdS ke dalam media tidak mempunyai kesan yang nyata terhadap jumlah tunas yang dihasilkan.

Terdapat hubung kait secara negatif antara tinggi tunas dan jumlah tunas yang dihasilkan. Tunas paling tinggi diperolehi pada media 1 mg/l and 2 mg/l BAP, sementara tunas paling pendek pada media 16 mg/l and 8 mg/l BAP. Kebalikan dari tinggi tunas, terdapat hubungan yang positif antara berat tunas dan jumlah tunas, dengan demikian media 16 mg/l and 8 mg/l BAP memberikan berat tunas yang tertinggi.

Selama subkultur, 2 mg/l BAP selalunya memberikan jumlah tunas yang paling banyak pada setiap subkultur (39.56 tunas per eksplan), diikuti oleh 1 mg/l BAP (27.71 tunas per eksplan). Secara am, tinggi tunas pada semua media subkultur tidak ada perbezaan yang bererti. Gabungan antara 1 mg/l BAP dan 2 mg/l Kinetin adalah merupakan media yang paling pantas untuk pemanjangan tunas (1.06 cm).

Media  $\frac{1}{2}$  MS + 15 mg/l NAA + 0.05% 'Charcoal' + 10 g/l 'sucrose' merupakan antara media yang paling sesuai untuk pemunculan akar, di mana jumlah akar yang tumbuh pada media ini mencapai 3.4 per tunas. Sedangkan media  $\frac{1}{2}$  MS +

1 mg/l IAA + 0.05% 'Charcoal' + 10 g/l 'sucrose' didapati paling sesuai untuk pemanjangan akar. Purata panjang akar pada media ini adalah 6.85 cm per tunas. Eksplan-eksplan dari pada kajian perakaran yang diaklimatisasi pada media tanah dicampur pasir dan gambut dengan nisbah 2 : 1: 1 dapat tumbuh dengan baik. 100% plantlet hidup pada kebuk kalbus (misting chamber) selama 8 minggu.

Dalam kajian organogenesis melalui induksi kalus, semua media kecuali MS, BAP sahaja dan 2,4-D sahaja dapat menghasilkan kalus.

Diharapkan, hasil daripada kajian ini dapat mempertingkatkan ilmu pengetahuan sains, khasnya pembiakan dengan kultur tisu, dan dapat diguna pakai sebagai rujukan sama ada oleh pelajar, penyelidik, mahupun pihak industri pertanian dalam penghasilan manggis secara besar-besaran.

## ACKNOWLEDGEMENTS

First, and most importantly, I am obligated to express gratitude to Allah, Most Merciful, for the blessing me granted onto me to execute and complete the study. I am very indebted to those who have helped me in preparing and conducting the study. I wish to express my deepest gratitude.

I would like to thank the Chairman of the Supervisory Committee, Associate Professor Mihdzar Abdul Kadir, PhD for his guidance and advice in the preparation of project proposal, implementation of research, presentation of seminars and completion of the thesis. I appreciate his patience and sincere approach to motivate, help, advise and guide me to finish my study at the Universiti Putra Malaysia. Thanks also are extended to my Committee members, Associate Professor Maheran Abdul Aziz, PhD in particular for her advice, comments, guidance, critical discussion and encouragement in the completion of this thesis and to Mr. Azmi Abdul Rashid who supported me by his encouragement and by the stimulating discussions held with me. They have given me the experience and knowledge, resulting in significant and positive impact on my work.

I am also deeply grateful to the Head of the Agency for Agricultural Research and Development, Department of Agriculture, Indonesia for awarding the scholarship that enables me to pursue a Master program at Universiti Putra

Malaysia, and to the Director of the Indonesia Centre Agricultural Agency for Biotechnology and Genetic Resources Development, Bogor, Indonesia for granting the study leave.

My appreciation goes to all staff of the Tissue Culture Laboratory, Department of Agriculture Technology, Universiti Putra Malaysia: Encik Rozaidi, Encik Ahmad Nazri, Cik Rostina Rolon, Cik Suharni, and Cik Ramisah for their help and assistance while conducting the study. I also wish to extend my gratefulness to Encik Rustam for his kind help to get the fresh-mangosteen fruits as study materials at the Agriculture Training Center in Kalumpang, Selangor - Darul Ehsan, Malaysia. I am indebted to Encik Zainuri, head the Department of Agriculture in Serdang, Selangor-Malaysia.for his help to get information on mangosteen in Malaysia. I am also indebted to Encik Mohammad Yusof Hashim, head of Fruits Division, Department of Agriculture in Kuala Lumpur, Malaysia.for the interest and kindness extended to me by asking Encik Anwar and Encik Amir Hamzah, staff of the Agricultural Department in Rembau, Negeri Sembilan - Darul Khusus and Agriculture Training Center in Kalumpang, Selangor - Darul Ehsan, Malaysia respectively, to provide me with fresh-mangosteen fruits as study material from their orchards. Without their assistance I would have difficulties conducting the experiments.

My special thanks are addressed to my housemates, Rozliza Ismail, Wan Rozidayani Wan Mohammad, Uswati Mohammad Ghazali, Norhazlina Nodin and Siti Khairani Khalid for sharing the joy and sorrow during my stay in Malaysia.

A special note of thanks also to Ibu Dr. Ir. Ika Mariska for her valuable encouragement and attention. Last but not least, I would like to express my deepest thanks to my father, Gito Soewarno and mother, Subariah for their support, encouragement and endless prayers. To my brothers, Teguh Setiono and Kukuh Setiyadi and my Sister, Sari Hening Andajani, thanks for your prayers and moral support. May God bless us all.

I certify that an Examination Committee has met on 25 April 2007 to conduct the final examination of Suci Rahayu on her Master of Science entitled “Micropropagation of Mangosteen (*Garcinia mangostana* L.) by Direct and Indirect Organogenesis” in accordance with Universiti Putra Malaysia (Higher Degree) Act 1980 and Universiti Putra Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Associate Professor Halimi Mohd Saud, PhD**

Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairperson)

**Associate Professor Saleh Kadzimin, PhD**

Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Associate Professor Datin Siti Nor Akmar Abdullah, PhD**

Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Ahmad Tarmidzi Hashim, PhD**

Malaysian Oil Palm Board  
(External Examiner)

---

**HASANAH MOHD. GHAZALI, PhD**

Professor/Deputy Dean  
School of Graduate Studies  
University Putra Malaysia

Date: 3 AUGUST 2007

This thesis submitted to Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

**Mihdzar Abdul Kadir, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Maheran Abdul Aziz, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Azmi Abdul Rashid, M.Phil**

Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

---

**AINI IDERIS, Ph.D**

Professor/Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 9 AUGUST 2007



## **DECLARATION**

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

---

**SUCI RAHAYU**

Date: 25 JUNE 2007



## TABLE OF CONTENTS

|  | <b>Page</b> |
|--|-------------|
| <b>DEDICATION</b>                                  | ii          |
| <b>ABSTRACT</b>                                    | iii         |
| <b>ABSTRAK</b>                                     | vi          |
| <b>ACKNOWLEDGEMENTS</b>                            | ix          |
| <b>APPROVAL</b>                                    | xii         |
| <b>DECLARATION</b>                                 | xiv         |
| <b>LIST OF TABLES</b>                              | xvii        |
| <b>LIST OF FIGURES</b>                             | xx          |
| <b>LIST OF PLATES</b>                              | xxii        |
| <b>LIST OF APPENDICES</b>                          | xxiii       |
| <b>LIST OF ABBREVIATIONS/NOTATIONS</b>             | xxix        |
| <br><b>CHAPTER</b>                                 |             |
| <br><b>1 GENERAL INTRODUCTION</b>                  | <br>1       |
| 1.1 Background                                     | 1           |
| 1.2 Objectives of the Study                        | 3           |
| <br><b>2 LITERATURE REVIEWS</b>                    | <br>4       |
| 2.1 Organogenesis                                  | 4           |
| 2.1.1 Direct Organogenesis                         | 5           |
| 2.1.2 Indirect Organogenesis                       | 8           |
| 2.2 Micropropagation Status of Mangosteen          | 14          |
| 2.2.1 Direct Organogenesis in Mangosteen culture   | 14          |
| 2.2.2 Indirect Organogenesis in Mangosteen Culture | 19          |
| 2.2.3 Acclimatization of Mangosteen Plantlet       | 20          |
| <br><b>3 DIRECT ORGANOGENESIS</b>                  | <br>22      |
| 3.1 Introduction                                   | 22          |
| 3.2 Study on Shoot Proliferation                   | 23          |
| 3.2.1 Materials and Methods                        | 23          |
| 3.2.2 Results                                      | 31          |
| 3.2.3 Discussions                                  | 79          |
| 3.3 Study on Shoot Elongation                      | 91          |
| 3.3.1 Materials and Methods                        | 91          |
| 3.3.2 Results                                      | 94          |

|          |                               |            |
|----------|-------------------------------|------------|
| 3.3.3    | Discussions                   | 102        |
| 3.4      | Study of Rooting              | 103        |
| 3.3.1    | Materials and Methods         | 104        |
| 3.3.2    | Results                       | 107        |
| 3.3.3    | Discussions                   | 114        |
| 3.5      | Study on Acclimatization      | 118        |
| 3.5.1    | Materials and Methods         | 118        |
| 3.5.2    | Results                       | 119        |
| 3.5.3    | Discussions                   | 121        |
| <b>4</b> | <b>INDIRECT ORGANOGENESIS</b> | <b>123</b> |
| 5.1      | Introduction                  | 123        |
| 5.2      | Materials and Methods         | 123        |
| 5.3      | Results                       | 130        |
| 5.4      | Discussions                   | 134        |
| <b>5</b> | <b>CONCLUSIONS</b>            | <b>137</b> |
|          | <b>REFERENCES</b>             | <b>140</b> |
|          | <b>APPENDICES</b>             | <b>154</b> |
|          | <b>BIODATA OF THE AUTHOR</b>  | <b>186</b> |

## LIST OF TABLES

| Table |   | Page |
|-------|---|------|
| 1     | The number of shoots produced per half seed explant of mangosteen on various proliferation media at first week of culture                             | 33   |
| 2     | The number of shoots produced per half seed explant of mangosteen on various proliferation media at second week of culture                            | 34   |
| 3     | The number of shoots produced per half seed explant of mangosteen on various proliferation media at third week of culture                             | 35   |
| 4     | The number of shoots produced per half seed explant of mangosteen on various proliferation media at fourth week of culture                            | 36   |
| 5     | Mean number of shoots produced per half seed explant of mangosteen on various proliferation media at fifth, sixth, seventh and eighth week of culture | 38   |
| 6     | The average shoot height produced from half seed explant of mangosteen on various proliferation media of culture at first week of culture             | 42   |
| 7     | The average shoot height produced from half seed explant of mangosteen on various proliferation media of culture at second week of culture            | 43   |
| 8     | The average shoot height produced from half seed explant of mangosteen on various proliferation media of culture at third week of culture             | 44   |
| 9     | The average shoot height produced from half seed explant of mangosteen on various proliferation media of culture at fourth week of culture            | 45   |
| 10    | The average shoot height produced from half seed explant of mangosteen on various proliferation media at fifth week of culture                        | 46   |

|    |   |    |
|----|---|----|
| 11 | The average shoot height produced from half seed explant of mangosteen on various proliferation media at sixth, seventh, and eighth week of culture | 47 |
| 12 | The weight of shoots produced per half seed explant of mangosteen on various proliferation media after eight weeks of culture                       | 53 |
| 13 | The number of shoots produced per half seed explant of mangosteen on various media at fourth week of subculture                                     | 58 |
| 14 | The number of shoots produced per half seed explant of mangosteen on various media at eighth week of subculture                                     | 59 |
| 15 | The number of shoots produced per half seed explant of mangosteen on various media at twelfth week of subculture                                    | 60 |
| 16 | The number of shoots produced per half seed explant of mangosteen on various media at sixteenth week of subculture                                  | 61 |
| 17 | The number of shoots produced per half seed explant of mangosteen on various media at twentieth week of subculture                                  | 62 |
| 18 | The average height of shoot produced from half seed explant of mangosteen on various media at fourth week of subculture                             | 67 |
| 19 | The average height of shoot produced from half seed explant of mangosteen on various media at eighth week of subculture                             | 68 |
| 20 | The average height of shoot produced from half seed explant of mangosteen on various media at twelfth week of subculture                            | 69 |
| 21 | The average height of shoot produced from half seed explant of mangosteen on various media at sixteenth week of subculture                          | 70 |

|    |  |     |
|----|--|-----|
| 22 | The average height of shoot produced from half seed explant of mangosteen on various media at twentieth week of subculture | 71  |
| 23 | The weight of shoots produced per half seed explant of mangosteen on various media at fourth week of subculture            | 75  |
| 24 | The weight of shoots produced per half seed explant of mangosteen on various media at eighth week of subculture            | 76  |
| 25 | The weight of shoots produced per half seed explant of mangosteen on various media at twelfth week of subculture           | 77  |
| 26 | The weight of shoots produced per half seed explant of mangosteen on various media at sixteenth week of subculture         | 78  |
| 27 | The weight of shoots produced per half seed explant of mangosteen on various media at twentieth week of subculture         | 79  |
| 28 | The average length of shoot on different types of elongation media   | 95  |
| 29 | The average weight of shoot on different types of elongation media   | 100 |
| 30 | Mean root number on mangosteen shoot cultured on various rooting media   | 108 |
| 31 | Mean root length on mangosteen shoot cultured on various rooting media   | 112 |
| 32 | The percentage of plantlet surviving under misting chamber for eight weeks   | 120 |
| 33 | Mean percentage of callus formation on media of BAP combined with concentration levels of 2,4-D                            | 131 |
| 34 | Mean percentage of callus formation on media of BAP combined with Glutamine  | 133 |

## LIST OF FIGURES

| Figure |   | Page |
|--------|---|------|
| 1      | Weekly mean number of shoots produced per half seed explant of mangosteen on media containing BAP and Kinetin either with or without AdS  | 32   |
| 2      | Weekly mean height of shoots produced per half seed explant of mangosteen on media containing BAP and Kinetin either with or without AdS  | 40   |
| 3      | The effect of BAP and Kinetin on the mangosteen shoot height at sixth, seventh and eighth week of culture   | 47   |
| 4      | The mean height of shoot on media containing BAP regardless of the AdS levels at sixth to eighth week of culture  | 48   |
| 5      | The mean height of shoot on media containing Kinetin regardless of the AdS levels at sixth to eighth week of culture  | 49   |
| 6      | Relationship between the height of shoot and the number of shoots produced per half seed explant of mangosteen at sixth, seventh and eighth week; $F = 27.19$ ; $df = 1, 16$ ; $P < 0.01$ )                       | 50   |
| 7      | Relationship between the initial fresh weight and the final fresh weight of mangosteen half seed explant; $F = 2.05$ ; $df = 1, 94$ ; $P > 0.05$  | 52   |
| 8      | Regression between the number of shoots produced per explant and the weight; $F = 69.63$ ; $df = 1, 22$ ; $P < 0.01$  | 54   |
| 9      | The number of shoots produced per explant, as affected by the interaction between AdS and medium of BAP and Kinetin at first to fifth four weekly of subculture. Means separation at each week by LSD at 5% level | 55   |
| 10     | The rate of increase in number of shoots produced from medium of 1 mg/l BAP, 2 mg/l BAP, and 4 mg/l BAP at first to fifth four weekly of subculture   | 57   |

|    |   |     |
|----|---|-----|
| 11 | Four weekly mean total height of shoots produced per half seed explant of mangosteen on media containing BAP and Kinetin either with or without AdS                         | 65  |
| 12 | The height of shoot on media enriched with AdS and without AdS at four weekly subculture  | 65  |
| 13 | The rate of increase in height of shoots produced on medium of 0-16 mg/l Kinetin at first to fifth four-weekly of subculture  | 72  |
| 14 | The growth of shoot explants on several media of elongation   | 97  |
| 15 | Relationship between the shoot height and the shoot weight at the final subculture (twenty four-week) of mangosteen shoot explant ( $F = 3.46$ ; $df = 1, 7$ ; $P > 0.05$ ) | 101 |
| 16 | The percentage of secondary root growing on the primary root of shoot explant on several rooting media after 28 weeks of subculture   | 109 |
| 17 | The relationship between concentration of 2,4-D and callus formation; $F = 3.42$ ; $df = 1, 3$ ; $P < 0.05$   | 132 |

## LIST OF PLATES

| Plate |   | Page |
|-------|---|------|
| 1     | The mangosteen. (A) Trees growing in the field, (B) Whole and half fruit, (C) Peeled seed before sterilization, and (D) Seed after sterilization  | 25   |
| 2     | Mangosteen shoots formed on media supplemented with 8 mg/l BAP (A) and 16 mg/l BAP (B) at eighth week of culture  | 39   |
| 3     | The shoot height attained on medium with 1 mg/l Kinetin (A) and 2 mg/l BAP (B) at eighth week of culture  | 49   |
| 4     | Mangosteen shoots formed on media with 1 mg/l BAP (A) and 2 mg/l BAP (B) at twentieth week of subculture  | 63   |
| 5     | Mangosteen shoots generated on media with 1 mg/l BA (A), 4 mg/l BA + 100 mg/l AdS (B), 4 mg/l Kinetin (C), and 4 mg/l Kinetin + 100 mg/l AdS (D) at twentieth week of subculture  | 73   |
| 6     | Shoot of mangosteen on elongation media with (A) 1 mg BAP + 2 mg/l Kinetin, (B) 0.5 mg/l BAP + 2 mg/l Kinetin, (C) 0.5 mg/l BAP + 4 mg/l Kinetin, and (D) MS without plant growth regulator at twenty fourth week of subculture | 98   |
| 7.    | The secondary root growing on the primary root of shoot explant on media with 10 mg/l IBA (A) and 5 mg/l NAA (B)  | 110  |
| 8.    | Root developed from shoot explants cultured on various rooting media. (A). IBA 1 mg/l, (B). IAA 1 mg/l, (C). NAA 0.5 mg/l, (D). $\frac{1}{2}$ MS 0, (E). NAA 10 mg/l, (F). NAA 15 mg/l  | 113  |
| 9.    | The eight weeks old plantlets growing in misting chamber  | 120  |
| 10.   | Callus formation on media with 5 mg/l 2,4-D + 1 mg/l BAP (A) and 10 mg/l 2,4-D + 2 mg/l BAP (B) on eight week of culture  | 133  |



## LIST OF APPENDICES

| Appendix<br>Table |  | Page |
|-------------------|--|------|
| 1                 | Analysis of variance for the number of shoots produced per half seed explant of mangosteen on various proliferation media at first week of culture         | 155  |
| 2                 | Analysis of variance for the number of shoots produced per half seed explant of mangosteen on various proliferation media at second week of culture        | 155  |
| 3                 | Analysis of variance for the number of shoots produced per half seed explant of mangosteen on various proliferation media at third week of culture         | 156  |
| 4                 | Analysis of variance for the number of shoots produced per half seed explant of mangosteen on various proliferation media at fourth week of culture        | 156  |
| 5                 | Analysis of variance for the number of shoots produced per half seed explant of mangosteen on various proliferation media at fifth week of culture         | 157  |
| 6                 | Analysis of variance for the number of shoots produced per half seed explant of mangosteen on various proliferation media at sixth week of culture         | 157  |
| 7                 | Analysis of variance for the number of shoots produced per half seed explant of mangosteen on various proliferation media at seventh week of culture       | 158  |
| 8                 | Analysis of variance for the number of shoots produced per half seed explant of mangosteen on various proliferation media at eighth week of culture        | 158  |
| 9                 | Analysis of variance for the average height of shoot produced from half seed explant of mangosteen on various proliferation media at first week of culture | 159  |